

Planar tropical gravitational descendants with one tangency condition

Falko Gauss

September 25, 2015

Abstract: We prove a formula which allows us to recursively compute planar tropical gravitational descendants which involve psi-classes of arbitrary power at marked ends fixed by points and additionally a psi-class of power one at exactly one marked end fixed by a line. Up to now almost exclusively tropical gravitational descendants with psi-classes at marked ends restricted by points had been considered.

Keywords: Tropical geometry, Gromov-Witten invariants, tropical intersection theory

2000 Mathematics Subject Classification: 14N35, 52B20

1 Introduction

Gromov-Witten theory is an important tool to tackle enumerative problems in algebraic geometry. The main objects of Gromov-Witten theory are the so-called Gromov-Witten invariants. They count (virtually) the number of curves of given genus and degree that pass through a certain number of chosen subvarieties. To also count curves that fulfill some tangency conditions, geometers started to use gravitational descendants which are generalized versions of Gromov-Witten invariants. Note that some authors also call these invariants descendant Gromov-Witten invariants (e.g. [7]).

With the emergence of tropical geometry also tropical counterparts of Gromov-Witten invariants and gravitational descendants were introduced. Unfortunately, the tropical gravitational descendants do not coincide with the corresponding classical numbers in general. This difference has structural reasons. The moduli space of marked tropical curves in \mathbb{R}^2 is non-compact and, hence, does not parameterize curves with components that approach the "boundary" of \mathbb{R}^2 in form of a limiting process. In particular, this fact makes the computation, which relies classically on the recursive structure of the boundary of the respective moduli space, much harder in the tropical setting. Nevertheless, some results on the computation of certain tropical gravitational descendants have been achieved. Most notably, Johannes Rau and Hannah Markwig proved

a tropical analogue to the WDVV Equation for planar invariants where psi-classes appear only at marked ends restricted by codimension two objects (see [7]). Here we want to go one step further and prove a Topological Recursion Relation which allows us to compute invariants which can additionally have a psi-class of power one at exactly one marked end restricted by a codimension one object. In order to do so we have to recall some notions from [3] and [10]. This will be done in Section 2 of the article at hand. Lastly, in Section 3 we will prove the desired formula using methods from [7] and [10].

For the sake of completeness, one should mention that there is very recent work of Peter Overholser which deals also with the computation of tropical descendant invariants in the plane (see [8]). But his point of view is more mirror symmetric and less intersection theoretic.

I would like to thank Andreas Gathmann for several helpful discussions. Note that this article contains material from my master thesis.

2 Defining tropical gravitational descendants

We consider the *moduli space* $\mathcal{M}_{0,l+m+n,trop}^{lab}(\mathbb{R}^2, d)$ of *rational parameterized tropical curves* in \mathbb{R}^2 of degree Δ with $l + m + n + |\Delta|$ labelled ends of which $l + m + n$ are contracted (see [3, Definition 4.1] for details). In analogy to the moduli space of pointed stable curves the contracted ends are called *marked ends*. In particular, we assume here that the degree is of the form $\Delta = \underbrace{\{(-e_1, -e_2, e_1 + e_2), \dots, (-e_1, -e_2, e_1 + e_2)\}}_{d \times}$, where e_1 and e_2 denote the unit

vectors in the plane \mathbb{R}^2 . By abuse of notation we will not distinguish between the labels of non-contracted ends and the corresponding elements in Δ . Furthermore, we will refer to $-e_1$, $-e_2$ and $e_1 + e_2$ as the *standard directions*.

Exemplarily, an element $(\Gamma, x_1, \dots, x_5, h)$ from $\mathcal{M}_{0,5,trop}^{lab}(\mathbb{R}^2, 2)$, i.e. a *marked tropical curve*, is shown in Figure 1.

According to what one sees Γ is an *abstract tropical curve*, i.e. a tree with a distinguished number of marked ends whose bounded edges are equipped with strictly positive lengths. The marked ends of the abstract tropical curve are visualized as dashed lines. The map $h : \Gamma \rightarrow \mathbb{R}^2$ maps Γ to an image as on the bottom of Figure 1. The image satisfies the balancing condition and marked ends are contracted to a point.

Now, we want to count marked tropical curves that fulfill certain incidence and tangency restrictions. Since we are working in the plane, we only have to consider incidence restrictions coming from codimension one objects, i.e. lines, and codimension two objects, i.e. points. The codimension zero object, i.e. the whole space \mathbb{R}^2 , does not give us non-trivial restrictions. So let G_j be certain (tropical) lines and P_k be certain points in \mathbb{R}^2 . For us a tropical line is a 3-valent graph with one vertex and one end in each of the standard directions (cf. the left-most curve in [7, Definition 2.7]). We will call the unique vertex of the




Figure 1: 5-marked tropical curve

tropical line its *root*.

We can partition the set of marked ends of a curve into three disjoint sets. In the following we will consider curves with $l + m + n$ marked ends and a partition $L \cup M \cup N = \{1, \dots, l + m + n\}$ with $|L| = l$, $|M| = m$ and $|N| = n$. We assume that the ends with labels $i \in L$ are unrestricted, the ends with labels $j \in M$ have to meet the respective lines G_j and the ends with labels $k \in N$ have to meet the respective points P_k . To make the meaning of the word "meet" precise, we need the following definition.

Definition 2.1. For $i = 1, \dots, l + m + n$ the map

$$ev_i : \mathcal{M}_{0, l+m+n, trop}^{lab}(\mathbb{R}^2, d) \rightarrow \mathbb{R}^2, (\Gamma, x_1, \dots, x_{l+m+n}, h) \mapsto h(x_i)$$

is called the *i-th evaluation map*.

By [3, Proposition 4.7] the moduli space of rational parameterized tropical curves can be equipped with a tropical (=weighted, balanced and polyhedral) fan structure via choosing one of the marked ends as the *anchor vertex*. So the evaluation maps become morphisms of fans. Moreover, in this sense we can talk about *cones*, *facets*, (*relative*) *interiors* and *rational functions* (cf. [7, Section 2]). Lastly, this enables us to apply the tropical intersection theory developed in [1] and to form *intersection products* consisting of rational functions on the fan $\mathcal{M}_{0, l+m+n, trop}^{lab}(\mathbb{R}^2, d)$. These intersection products have the structure of

tropical complexes and, hence, we can talk about *polyhedra* and the *dimension* of an intersection product.

Now we want to enrich these intersection products further by allowing *psi-classes* ψ_i at the marked ends with indices $i \in L \cup M \cup N$ (see [7, Definition 2.2] for the definition of a psi-class in tropical geometry). They can be thought of as encoding tangency (or higher order contacts) of the curves at the respective marked ends. In formal terms a psi-class ψ_i is the codimension one subfan of $\mathcal{M}_{0,l+m+n,trop}^{lab}(\mathbb{R}^2, d)$ that consists of cones corresponding to trees where the marked end i sits at a vertex of valence ≥ 4 . Let t_i , s_j and r_k be the numbers that specify how many psi-classes at $i \in L$, $j \in M$ and $k \in N$, respectively, are required.

Definition 2.2 (cf. [7, Proposition and Definition 2.10]). Let d, l, m, n, t_i, s_j and r_k be non-negative integers which satisfy:

$$l + m + n + 3d - 3 + 2 = m + 2n + \sum_{i \in L} t_i + \sum_{j \in M} s_j + \sum_{k \in N} r_k. \quad (1)$$

Then the respective *tropical gravitational descendant* is defined as:

$$\left\langle \prod_{i \in L} \tau_{t_i}(0) \prod_{j \in M} \tau_{s_j}(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d := \frac{1}{(d!)^3} \deg \left(\prod_{i \in L} \psi_i^{t_i} \prod_{j \in M} ev_j^*(G_j) \psi_j^{s_j} \prod_{k \in N} ev_k^*(P_k) \psi_k^{r_k} \cdot \mathcal{M}_{0,l+m+n,trop}^{lab}(\mathbb{R}^2, d) \right).$$

Note that the dimension of $\mathcal{M}_{0,l+m+n,trop}^{lab}(\mathbb{R}^2, d)$ is $l + m + n + 3d - 3 + 2$ by [3, Proposition 4.7] and the codimension of the intersection of psi-classes is $\sum_{i \in L} t_i + \sum_{j \in M} s_j + \sum_{k \in N} r_k$ where the pullback of a line has codimension 1 and the pullback of a point codimension 2. So the lengthy Equation 1 enforces the intersection product to be zero-dimensional. This assumption is necessary to talk about the *degree* of an intersection product. The definition above is independent of the choice of the lines G_j and the points P_k (cf. Section 2 of [7]). In particular, the definition also makes sense if we choose rather special configurations of lines and points. But to give some enumerative meaning to the definition we have to assume a certain generality of the lines G_j and the points P_k . Namely, we assume that the lines G_j and the points P_k are in *general position* in the sense of [3, Definition 3.2]. It is worth mentioning that the assumption that G_j and P_k are in general position is not a serious limitation because the P_k and G_j in general position form a dense subset in the set of P_k and G_j under all conditions (which can be identified with some large-dimensional real space). A proof of this fact is presented in [7, Lemma 3.4].

Assuming that the lines G_j and points P_k are in general position, the intersection product consists (in set-theoretic terms) of the tropical curves that pass

through G_j and P_k equipped with certain weights. The degree of the intersection product, to wit the tropical gravitational descendant, is then the sum of all weights. According to [2, Lemma 2.4] (or [9, Remark 2.2.14]) these weights can be computed as the product of the weight of the facet the respective curve lies in and some determinantal expression. Nevertheless, the computation of the weights via determinantal formulas is very unwieldy and unsuitable for a large degree or a large number of marked ends. That is why one wish for recursion formulas, like for instance the Topological Recursion Relation in classical Gromov-Witten theory. Such formulas rely classically on the well-known Splitting Lemma (e.g. see [6, 5.2.1 Lemma]). Johannes Rau developed a tropical version of the Splitting Lemma. However, his Tropical Splitting Lemma [10, Theorem 4.13] only works under rather strict assumptions. In fact, this is one of the major problems when transferring the classical Gromov-Witten theory to the tropical setting. An example where the Tropical Splitting Lemma fails can be found in [10, Remark 4.18]. Here we restrict ourselves to a special situation.

Assumption 2.3. From now on (unless stated otherwise) we assume that $L = \emptyset$, $M = \{1\}, \{2, 3\} \subseteq N$ and $s_1 = 1$.

So we have a fan $X := \psi_1^0 \prod_{k \in N} \psi_k^{r_k} \cdot \mathcal{M}_{0,1+n,trop}^{lab}(\mathbb{R}^2, d)$ and a one-dimensional intersection product $F := (\tau_0(1) \prod_{k \in N} \tau_{r_k}(2))$. This situation has two major advantages. Firstly, the weight $w_X(\sigma)$ of the facet σ the respective curve lies in is one by [7, Theorem 5.3] and, therefore, can be omitted in this article. Secondly, we can apply the Tropical Splitting Lemma by [10, Lemma 4.16]. The Tropical Splitting Lemma splits the curves of an intersection product at their *contracted bounded edges* as described in [4, Remark 2.7]. Note that a contracted bounded edge induces a *reducible partition* $I|J$ of the labels of the marked ends of the curve, i.e. a partition into two non-empty sets I and J such that directions in $I \cap \Delta$ (and hence also in $J \cap \Delta$) sum up to the zero vector. In our situation the Tropical Splitting Lemma [10, Theorem 4.13] takes the form of:

$$\begin{aligned}
& \left\langle \tau_1(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d = \\
& \sum_{\substack{I|J \text{ reducible} \\ 1 \in I, 2, 3 \in J}} \sum_{\varepsilon + \zeta = 2} \left\langle \tau_0(\varepsilon) \tau_0(1) \prod_{k \in N \cap I} \tau_{r_k}(2) \right\rangle_{d_1} \cdot \left\langle \tau_0(\zeta) \prod_{k \in N \cap J} \tau_{r_k}(2) \right\rangle_{d_2} \\
& + \left\langle \phi_{1|2,3} \tau_0(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d. \tag{2}
\end{aligned}$$

In Equation (2) the rational function $\phi_{1|2,3}$ is defined as specified in [10, Section 5.2]. Here, it is sufficient for us to know that $\phi_{1|2,3}$ is of the following form (see [10, Lemma 5.6]) on each facet σ of F , i.e. on each top-dimensional

polyhedron of F :

$$\phi_{1|2,3}|_{\sigma} = \begin{cases} \text{constant,} & \text{if the interior curves of } \sigma \text{ have} \\ & \text{a contracted bounded edge} \\ \text{sum of lengths of edges that} & \\ \text{separate 1 from 2, 3,} & \text{otherwise.} \end{cases} \quad (3)$$

Loosely speaking $\phi_{1|2,3}$ decides if the result applying the Tropical Splitting Lemma equals the result one obtains by applying the classical Splitting Lemma to the respective classical invariants. This is the case in almost all of the existing literature (e.g. in [7] and [10]), because there $\phi_{1|2,3}$ is bounded. Here $\phi_{1|2,3}$ shows a slightly different behaviour. This will be made precise in the next corollary.

Definition 2.4. Let C be a curve in $\mathcal{M}_{0,1+n,trop}^{lab}(\mathbb{R}^2, d)$. A subgraph \mathcal{S} of C that is homeomorphic to \mathbb{R} and does not involve (the closure of) a marked end x_k with $k \in N$ is called a *string*.

Corollary 2.5. Let F be as above. Then, the function $\phi_{1|2,3}$ can be unbounded only on facets of F with interior curves that look like in Figure 2 (where E and G_1 can run along any of the three standard directions).




Figure 2: Deformation which yields an unbounded $\phi_{1|2,3}$

Proof. Every facet of F is described by the deformation of one of its interior curves, because facets are one-dimensional. A list of all possible deformations is presented in [10, Lemma 5.13]. In our situation ($L = \emptyset$) the list looks as follows. The deformation of an interior curve C in a facet σ of F is described by one of the following cases:

1. C contains a contracted bounded edge and one can vary its length arbitrarily.
2. C does not have a contracted bounded edge; and
 - (a) C has a *degenerated* vertex V of valence three, i.e. a vertex of one of the following types:
 - i. The vertex is incident to a bounded edge E , a non-contracted end and the marked end with label 1 such that the curve C and the line G_1 do not intersect transversally at $ev_1(C)$.

- ii. The vertex is incident to a bounded edge E and two non-contracted ends such that the span of the non-contracted ends near V is still only one-dimensional.

The deformation is given by changing the length of the bounded edge E .

- (b) C contains a string \mathcal{S} with two non-contracted ends and only one adjacent bounded edge E as shown in Figure 3 such that all vertices of \mathcal{S} are non-degenerated and of valence three in C . This string can be moved by changing the length of E .

Now we will work through this list and stick to the notations from there. Assume that σ is a facet of F which has an interior curve with a contracted bounded edge, i.e. σ looks like 1. on the list. Then $\phi_{1|2,3}$ is constant on σ (3) and, thus, bounded on σ .

Now, assume that σ is a facet with an interior curve as in 2. (a) on the list. As $\phi_{1|2,3}$ measures the sum of the lengths of the edges that separate 1 from 2, 3, the function $\phi_{1|2,3}$ can only be unbounded on σ if it takes the length of the growing edge E into account. Note that, in the case 2. (a) ii. E does not separate $1 \in M$ from 2, 3 and, hence, does not contribute to $\phi_{1|2,3}$ on σ . This holds, because the vertex V in 2. (a) ii. is of valence three and, hence, cannot also be incident to the marked end with label 1. In the case 2. (a) i. the only chance that E is taken into account by $\phi_{1|2,3}$ is the situation shown in Figure 2. Finally, assume that σ is a facet as in 2. (b) on the list. In this case an interior curve of σ looks like in Figure 3. Namely, E only contributes to $\phi_{1|2,3}$ on σ , if E separates 1 from 2, 3. So by construction x_1 has to lie on \mathcal{S} (cf. [9, Lemma 2.4.14]) and x_2, x_3 have to lie "behind" E . Moreover, on such a curve E must have one of the standard directions (i.e. directions of the edges of G_1), because otherwise the edge E will not grow infinitely for obvious reasons. But the line G_1 was chosen to look like the left-most curve in [7, Definition 2.7]. Hence, it has a root and none of the half-rays can be infinitely long in both directions. In Figure 3 this means that the line G_1 on the left hand side has to split somewhere, i.e. the root has to appear. So E cannot grow infinitely and the situation in Figure 3 cannot cause an unbounded $\phi_{1|2,3}$ either.

All in all, the only interior curves which can cause facets with unbounded $\phi_{1|2,3}$ are the ones in Figure 2. \square

3 A new Topological Recursion Relation

A very helpful technique to compute certain intersection products, resp. their degrees, is (what we will call) the λ -method. The λ -method is due to Andreas Gathmann and Hannah Markwig (see [4]). To apply the λ -method one adds a factor $ft^*(\lambda)$ to a one-dimensional intersection product of interest. This factor is the pull-back of an element $\lambda \in \mathcal{M}_{0,4,trop}$ under the forgetful map $ft : \mathcal{M}_{0,l+m+n,trop}^{lab}(\mathbb{R}^2, d) \rightarrow \mathcal{M}_{0,4,trop}$ which forgets all ends except those with




Figure 3: Bounded deformation of a string

the labels 1, 2, 3, 4 of a given curve. Here $\mathcal{M}_{0,4,trop}$ is the *moduli space of 4-marked abstract tropical curves* as defined in [3, Definition 3.2]. This moduli space consists of four different types of curves. All different types are shown in [4, Example 1.1]. In the following we omit the case that λ is just the single vertex in $\mathcal{M}_{0,4,trop}$, i.e. looks like the right-most curve in [4, Example 1.1]. Then, by abuse of notation, λ denotes not only an element in $\mathcal{M}_{0,4,trop}$ but also a real number. This real number is the length of the unique bounded edge (cf. [4, Example 1.1]). Along these lines, the forgetful map ft can be considered as a map to the set of real numbers.

Curves in an intersection product that involves $ft^*(\lambda)$ fulfill the additional requirement that they map to λ under ft . Now, the key idea of the λ -method is to make λ sufficiently large. Then, the curves in the intersection product have certain special properties, but the degree of the intersection product remains the same by the so-called rational equivalence (cf. the proof of [7, Theorem 8.1]).

The overall goal of this section is to recursively compute invariants of the form $\left\langle \tau_0(0)^{l'} \tau_1(0)^{l''} \tau_0(1)^{m-1} \tau_1(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d$ where $l', l'' \in \mathbb{N}$ and $m \in \mathbb{N}_{>0}$. We will see that in order to do so, it is sufficient to derive a recursion formula for invariants of the form $\left\langle \tau_1(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle$, i.e. invariants as they appeared in Section 2. As we learned in the last section the curves shown in Figure 2 are of particular importance. The next lemma is the essential step to describe these curves in the realm of the λ -method.

Lemma 3.1. *Let F be as in the previous section and σ be a facet of F . Moreover, let C be an interior curve of σ as shown in Figure 2, such that x_4 with*

$4 \in \Delta$ is the unbounded edge adjacent to the growing bounded edge E and C maps to λ under ft . Then the linear parts $(\phi_{1|2,3})_\sigma$ and $(ft)_\sigma$ of the restricted functions $\phi_{1|2,3}|_\sigma$ and $ft|_\sigma$ coincide.

Proof. We know from [10, Lemma 5.13] that we get other curves of σ by varying the length of the bounded edge E of C . As σ is one-dimensional all curves in σ arise in that way. In particular, all interior curves of σ do not have a contracted bounded edge and we can use the second description of $\phi_{1|2,3}$ from Equation (3). Let \tilde{C} be a curve in σ and \tilde{E} be the deformed edge which originates from E , then we obtain:

$$\begin{aligned} \phi_{1|2,3}(\tilde{C}) &=_{(3)} \text{sum of lengths of edges that separate 1 from 2, 3} \\ &= \text{length of } \tilde{E} + \text{term that is constant on } \sigma, \end{aligned}$$

On the other hand we obtain:

$$ft(\tilde{C}) = \text{length of } \tilde{E}.$$

Since \tilde{C} was an arbitrary curve in σ , we can conclude that $\phi_{1|2,3}|_\sigma + \text{constant} = ft|_\sigma$. So the linear parts of $\phi_{1|2,3}|_\sigma$ and $ft|_\sigma$ coincide. \square

Now we have to quantify the weight of curves as shown in Figure 2. As already mentioned such weights are given by determinantal expressions. A nice example of how to compute such an absolute determinant $|\det_C(\dots)|$ for a curve C is presented in [2, Example 2.6]. However, here we use notions analogous to [7, Notation 5.2]. In particular, we consider evaluation maps

$$ev : X \rightarrow \mathbb{R}^1 \times (\mathbb{R}^2)^n, C \mapsto ((ev_1(C))_x, (ev_k(C))_{k \in N}), \quad (4)$$

$$\tilde{ev} : X \rightarrow (\mathbb{R}^2)^{n+1}, C \mapsto ((ev_1(C)), (ev_k(C))_{k \in N}), \quad (5)$$

where the index x at a point in \mathbb{R}^2 indicates its first coordinate. Later on we will also use the index y to indicate a second coordinate. So the evaluation map \tilde{ev} arises from ev by replacing the line restriction at x_1 with a point restriction at x_1 .

Lemma 3.2. *Let $Z = (ft^*(\lambda) \tau_0(1) \prod_{k \in N} \tau_{r_k}(2))$ be a zero-dimensional intersection product and X be as above. Let $4 \in \Delta$. Moreover, let C be an interior curve in a facet of F as shown in Figure 2, such that x_4 is the unbounded edge adjacent to the growing bounded edge E and C maps to a large λ under ft . Then, the weight of C in Z is $|\det_C(ft \times ev)| = |\det_C(\tilde{ev})|$.*

Proof. Let C be a curve as above. Let l_1 be the length of its growing bounded edge E (see Figure 2). We want to compute the weight of C in Z . In so doing, we choose a marked end $x_a \neq x_1$ as the anchor vertex. Such a marked end must exist, because the number of marked ends is greater than three by Assumption

2.3. The position of x_a is determined by the coordinates $(h(x_a))_x$ and $(h(x_a))_y$. Say E runs along the standard direction $-e_1$, then the position of the marked end x_1 is given as

$$h(x_1) = h(x_a) - l_1 \cdot e_1 + \underbrace{\quad \quad \quad}_{\text{expression in the other bounded lengths}}. \quad (6)$$

The descriptions of the positions of all other marked ends do not involve the length l_1 , because we chose $x_a \neq x_1$.

Now, by [2, Lemma 2.4] we know that the weight of C is given as the absolute value of the determinant $|\det_C(ft \times ev)|$. We call the underlying matrix of this absolute determinant H . Let the line G_1 be given by $\max\{y, c_1\}$. Thus, the pull-back of the evaluation map with plugged in coordinates from (6) is of the form $\max\{(h(x_a))_y + \dots - c_1, 0\}$. We assume that the points P_k are given by functions of the form $\max\{x, p_1^k\}$ and $\max\{y, p_2^k\}$. Then, the pull-backs of the evaluation maps at the points are of the form $\max\{(h(x_k))_x - p_1^k, 0\}$ and $\max\{(h(x_k))_y - p_2^k, 0\}$. So we can determine the entries of all rows of H that correspond to pull-backs of evaluations at the line G_1 and the points in the columns corresponding to l_1 , $(h(x_a))_x$ and $(h(x_a))_y$. The only missing row is the row corresponding to the pull-back of the $\mathcal{M}_{0,4,trop}$ -coordinate λ under ft . But, since λ is large and there is no contracted bounded edge in a curve in this facet, the length l_1 must contribute to the length of λ . This causes an entry one at the column corresponding to l_1 . All in all, the entries in H are summarized in the table below.

	$(h(x_a))_x$	$(h(x_a))_y$	l_1	other bounded lengths
evaluation at G_1	0	1	0	*
evaluation at $(P_k)_x$ for $k \in N$	1	0	0	*
evaluation at $(P_k)_y$ for $k \in N$	0	1	0	*
ft	0	0	1	*

We can permute rows and columns of H , because we are only interested in the absolute value of its determinant. So we permute the rows and columns of H in such a way that its first row is the row corresponding to ft and that this row starts with a one. Then, we can apply the determinantal formula for block matrices on the upper left 1 by 1 matrix block. This means we forget the row corresponding to ft and the column corresponding to l_1 . Now, applying the determinantal formula for block matrices backwards we can reintroduce a column corresponding to l_1 and introduce a row corresponding to $(P_1)_x$ for a new point P_1 which lies on the end pointing towards $-e_1$ of the line G_1 . We end up with the matrix entries shown in the table below.

	l_1	$(h(x_a))_x$	$(h(x_a))_y$	other bounded lengths
evaluation at $(P_1)_x$	-1	1	0	*
evaluation at $G_1 = (P_1)_y$	0	0	1	*
evaluation at $(P_k)_x$ for $k \in N$	0	1	0	*
evaluation at $(P_k)_y$ for $k \in N$	0	0	1	*

But this is exactly the matrix of the curve C for the evaluation map $\tilde{e}v$ which arises from ev by replacing the line restriction at x_1 with a point restriction given by P_1 . This yields $|\det_C(ft \times ev)| = |\det_C(\tilde{e}v)|$ and, hence, proves the claim. \square

Finally we have all tools available to prove the main result of this article.

Theorem 3.3. *It holds:*

$$\begin{aligned} & \left\langle \tau_1(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d = \\ & \sum \left\langle \tau_0(\varepsilon) \tau_0(1) \prod_{k \in N \cap I} \tau_{r_k}(2) \right\rangle_{d_1} \cdot \left\langle \tau_0(\zeta) \prod_{k \in N \cap J} \tau_{r_k}(2) \right\rangle_{d_2} \\ & + 3 \left\langle \tau_0(2) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d \end{aligned}$$

where the sum ranges over all $\varepsilon + \zeta = 2$ with $\varepsilon, \zeta \in \mathbb{Z}_{\geq 0}$ and $d_1 + d_2 = d$ with $d_1, d_2 \in \mathbb{Z}_{\geq 0}$ and reducible partitions $I|J$ with $1 \in I$ and $2, 3 \in J$.

Proof. By (2) we can write $\left\langle \tau_1(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d$ as

$$\begin{aligned} & \left\langle \tau_1(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d = \\ & \sum_{\substack{I|J \text{ reducible} \\ 1 \in I, 2, 3 \in J}} \sum_{\varepsilon + \zeta = 2} \left\langle \tau_0(\varepsilon) \tau_0(1) \prod_{k \in N \cap I} \tau_{r_k}(2) \right\rangle_{d_1} \cdot \left\langle \tau_0(\zeta) \prod_{k \in N \cap J} \tau_{r_k}(2) \right\rangle_{d_2} \\ & + \left\langle \phi_{1|2,3} \tau_0(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d. \end{aligned}$$

It remains to quantify $\left\langle \phi_{1|2,3} \tau_0(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d$. For reasons of clarity, we denote the set of facets of F as $F^{(1)}$ and the set of curves, to wit vertices, of F as $F^{(0)}$. Then by the definition of tropical divisors [1, Definition 3.4] it holds

$$\left\langle \phi_{1|2,3} \tau_0(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d = \sum_{F^{(0)} \ni C < \sigma \in F^{(1)}} w_F(\sigma) (\phi_{1|2,3})_{\sigma}(u_{\sigma/C}), \quad (7)$$

where $w_F(\sigma)$ is the weight of the facet σ in F and $u_{\sigma/C}$ is the primitive generator of the integer vector pointing from C to σ . For facets $\sigma \in F^{(1)}$ that contain two different vertices the two corresponding primitive generators have opposite directions (see [9, Lemma 1.4.4]). Thus the two respective summands in (7) cancel out. The $\sigma \in F^{(1)}$ that do not contain two different vertices are unbounded polyhedra. We know from Corollary 2.5 that $\phi_{1|2,3}$ is bounded on all facets of F which do not have interior curves like in Figure 2. But on unbounded polyhedra a rational function can only be bounded if it is constant, i.e. $(\phi_{1|2,3})_\sigma \equiv 0$. So the summands in (7) belonging to polyhedra which do not have interior curves like in Figure 2 vanish. In fact, it holds:

$$\left\langle \phi_{1|2,3}\tau_0(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d = \sum_{\substack{F^{(0)} \ni C < \sigma \in F^{(1)}: \\ \sigma \text{ has interior curve} \\ \text{like in Figure 2}}} w_F(\sigma) (\phi_{1|2,3})_\sigma (u_{\sigma/C}). \quad (8)$$

Due to Lemma 3.1 the linear parts $(\phi_{1|2,3})_\sigma$ and $(ft)_\sigma$ coincide. Thus, it follows

$$\left\langle \phi_{1|2,3}\tau_0(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d = \sum_{\substack{F^{(0)} \ni C < \sigma \in F^{(1)}: \\ \sigma \text{ has interior curve} \\ \text{like in Figure 2}}} w_F(\sigma) (ft)_\sigma (u_{\sigma/C}). \quad (9)$$

Again, by [2, Lemma 2.4] we can reformulate (9) and write (sticking to the notations from Lemma 3.2)

$$\left\langle \phi_{1|2,3}\tau_0(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d = \sum_{\substack{C \text{ looks like} \\ \text{in Figure 2}}} |\det_C(ft \times ev)|. \quad (10)$$

Here, as well as in the next two equations, the curves C over which we sum up live in the subcomplex containing the points of X that map to the correct point P_k for all $k \in N$ under the respective evaluation map and map to a fixed sufficiently large λ under ft . Now, we can divide the curves that look like in Figure 2 into three classes depending along which standard direction the end

x_4 , which is the unbounded edge adjacent to the marked end x_1 , runs. It holds:

$$\begin{aligned}
\left\langle \phi_{1|2,3} \tau_0(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d &= \sum_{\substack{C \text{ looks like} \\ \text{in Figure 2}}} |\det_C(ft \times ev)| \\
&= \sum_{\substack{C \text{ looks like in} \\ \text{Figure 2 and} \\ x_4 \text{ runs along } -e_1}} |\det_C(ft \times ev)| \\
&\quad + \sum_{\substack{C \text{ looks like in} \\ \text{Figure 2 and} \\ x_4 \text{ runs along } -e_2}} |\det_C(ft \times ev)| \\
&\quad + \sum_{\substack{C \text{ looks like in} \\ \text{Figure 2 and} \\ x_4 \text{ runs along } e_1 + e_2}} |\det_C(ft \times ev)|.
\end{aligned}$$

Finally, from Lemma 3.2 we obtain

$$\begin{aligned}
\left\langle \phi_{1|2,3} \tau_0(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d &= \sum_{\substack{C \text{ looks like in} \\ \text{Figure 2 and} \\ x_4 \text{ runs along } -e_1}} |\det_C(\tilde{e}v')| \\
&\quad + \sum_{\substack{C \text{ looks like in} \\ \text{Figure 2 and} \\ x_4 \text{ runs along } -e_2}} |\det_C(\tilde{e}v'')| \\
&\quad + \sum_{\substack{C \text{ looks like in} \\ \text{Figure 2 and} \\ x_4 \text{ runs along } e_1 + e_2}} |\det_C(\tilde{e}v''')|, \quad (11)
\end{aligned}$$

where, $\tilde{e}v'$, $\tilde{e}v''$ and $\tilde{e}v'''$ arise from ev by replacing the line restriction at x_1 with a point restriction given by new points $P_{1'}$, $P_{1''}$ and $P_{1'''}$ on G_1 (in the sense of Lemma 3.2). The points will be chosen in a specific way as described later in the case of $P_{1'}$. Now we observe that the sums in (11) look almost like the number (see [7, Theorem 5.3]) $\langle \tau_0(2) \prod_{k \in N} \tau_{r_k}(2) \rangle = \sum_{C \in S} |\det_C(\tilde{e}v)|$ where S is the subcomplex of X containing all points in X that map to the correct point P_k for $k \in N \cup \{1\}$ under the respective evaluation map. So it is natural to try to express each of the three sums via the invariant $\langle \tau_0(2) \prod_{k \in N} \tau_{r_k}(2) \rangle$. In so doing, we have to relate the index of summation of each of the sums in (11) to the index of summation occurring in the sum expression of $\langle \tau_0(2) \prod_{k \in N} \tau_{r_k}(2) \rangle$. Making this precise, we define three subcomplexes S_1 , S_2 and S_3 of X where S_1 contains all points in X that map to the correct point P_k for $k \in N \cup \{1'\}$ under the respective evaluation map, S_2 contains all points that map to P_k for $k \in N \cup \{1''\}$ and S_3 contains all points that map to P_k for $k \in N \cup \{1'''\}$. We want to show the following three equalities of (finite) sets:

$$\{C \mid C \text{ looks like in Figure 2 and } x_4 \text{ runs along } -e_1\} = S_1 \quad (12)$$

$$\{C \mid C \text{ looks like in Figure 2 and } x_4 \text{ runs along } -e_2\} = S_2 \quad (13)$$

$$\{C \mid C \text{ looks like in Figure 2 and } x_4 \text{ runs along } e_1 + e_2\} = S_3. \quad (14)$$

This then proves the claim, because by (11) we could write

$$\left\langle \phi_{1|2,3} \tau_0(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d = 3 \left\langle \tau_0(2) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d.$$

So let us check the two inclusions for S_1 . Subsequently, the argument can be carried out analogously for S_2 and S_3 .

First of all, we assume that the line G_1 and the points P_k for $k \in N \cup \{1'\}$ are organized in the following way:

- the x - and y -coordinates of the root of the line G_1 and of the points P_k for $k \in N$ are in an open interval $(-\epsilon, \epsilon)$ for some small real number $\epsilon > 0$;
- the point $P_{1'}$ lies on the end of the line G_1 which points into the direction $-e_1$ and its x -coordinate is $-\lambda$ where $\lambda \gg 0$ is some large real number;
- (additionally) the point P_2 lies at the origin and the x -coordinate of P_3 lies in the interval $[0, \epsilon]$.

This means we assume that the point $P_{1'}$ lies "far away" from the other points P_k for $k \in N$ (see Figure 4). Nevertheless, we still assume that all the points are in general position. This is possible because the points in general position are dense in the set of all points (cf. Section 2). Note that we are now in the situation of [11, Lemma 2.2.1]. As a consequence of this lemma we get the two desired inclusions.

" \subseteq ": Let C be a curve that looks like in Figure 2 and its unbounded edge x_4 runs along $-e_1$. Particularly, the edge x_4 runs along the end of G_1 pointing towards $-e_1$ and it is incident to a unique vertex. This vertex has an x -coordinate greater than $-\lambda$, because C maps to λ under ft by assumption and the points P_2 and P_3 have an x -coordinate in the interval $[0, \epsilon]$. Furthermore, the point $P_{1'}$ lies on the end of G_1 pointing towards $-e_1$ and it is placed at an x -coordinate of $-\lambda$ by construction. Thus, the unbounded edge x_4 of C has to meet $P_{1'}$. All in all, the curve C maps to $P_{1'}$ and P_k for $k \in N$ under the respective evaluation maps. So C is in S_1 .

" \supseteq ": Let $C \in S_1$. We want to show that C must look like in Figure 4, i.e. especially like in Figure 2. Therefore, assume that C does not look like in Figure 4. That means we assume that x_1 is not adjacent to an unbounded edge. By the balancing condition one of its adjacent bounded edges must be pointing leftwards and as this edge is bounded it is adjacent to a vertex with an x -coordinate smaller than $-\lambda$. But such a vertex does not exist, because all vertices of C must lie in $R := \{(x, y) \in \mathbb{R}^2 \mid -\lambda \leq x \leq \epsilon, -\epsilon \leq y \leq \epsilon\}$ by [11, Lemma 2.2.1]. So this case cannot occur. Hence, the curve $C \in S_1$ looks like in Figure 4 and the unbounded edge x_4 , which is the unbounded edge adjacent to x_1 , runs along $-e_1$. \square




Figure 4: Curve C in $(\tau_0(2) \prod_{k \in N} \tau_{r_k}(2))$ for special point configuration

By Theorem 3.3 we are able to compute all tropical gravitational descendants of the form $\langle \tau_1(1) \prod_{k \in N} \tau_{r_k}(2) \rangle_d$. Using in addition the Tropical String Equation [10, Theorem 3.12], the Tropical Divisor Equation [10, Theorem 3.17] and the Tropical Dilaton Equation [10, Theorem 3.13] we can recursively compute all tropical gravitational descendants of the form

$$\left\langle \tau_0(0)^{l'} \tau_1(0)^{l''} \tau_0(1)^{m-1} \tau_1(1) \prod_{k \in N} \tau_{r_k}(2) \right\rangle_d. \quad (15)$$

Remark 3.4. This means we are in particular able to compute the tropical counterparts of the so-called first descendant invariants with exactly one tangency condition $\tau_1(1)$. First descendant invariants are gravitational descendants which involve only psi-classes of power at most one. They play an important role in many classical enumerative questions because they are sufficient to study characteristic number problems (cf. [5]).

To close this article and to see the new recursion formula in action, we want to compute the invariant $\langle \tau_1(1) \tau_1(2) \tau_1(2) \rangle_2$. In fact, it holds:

$$\begin{aligned}
\langle \tau_1(1) \tau_1(2) \tau_1(2) \rangle_2 &=_{3.3} \left(\sum \langle \tau_0(\varepsilon) \tau_0(1) \rangle_{d_1} \cdot \langle \tau_0(\zeta) \tau_1(2) \tau_1(2) \rangle_{d_2} \right. \\
&\quad \left. + 3 \underbrace{\langle \tau_0(2) \tau_1(2) \tau_1(2) \rangle_2}_{= \tau_1^1} \right) \\
&= \underbrace{\langle \tau_0(0) \tau_0(1) \rangle_0}_{=0} \cdot \langle \tau_0(2) \tau_1(2) \tau_1(2) \rangle_2 + 3 \\
&= 3. \tag{16}
\end{aligned}$$

The numerical value of the algebro-geometric version $\langle \tau_1(1) \tau_1(2) \tau_1(2) \rangle_2^{alg}$ of this invariant equals 0 and can be easily computed via standard techniques (e.g. see [6]). So this is an example where the classical algebro-geometric and the tropical number do not coincide as mentioned in the introduction.

References

- [1] L. Allermann and J. Rau, *First Steps in Tropical Intersection Theory*, Math. Z. **264** (2010) Issue 3, 633–670.
- [2] M. Franz and H. Markwig, *Tropical enumerative invariants of \mathbb{F}_0 and \mathbb{F}_2* , Adv. Geom. **11** (2011) No. 1, 49–72.
- [3] A. Gathmann, M. Kerber and H. Markwig, *Tropical fans and the moduli spaces of tropical curves*, Compos. Math. **145** (2009) No. 1, 173–195.
- [4] A. Gathmann and H. Markwig, *Kontsevich’s formula and the WDVV equations in tropical geometry*, Adv. Math. **217** (2008) No. 2, 537–560.
- [5] T. Graber, J. Kock and R. Pandharipande, *Descendant invariants and characteristic numbers*, Amer. J. Math. **124** (2002) No. 3, 611–647.
- [6] J. Kock, *Notes on Psi classes*, available at <http://mat.uab.es/kock/GW/notes/psi-notes.pdf>.
- [7] H. Markwig and J. Rau, *Tropical descendant Gromov-Witten invariants*, Manuscripta Math. **129** (2009) Issue 3, 293–335.
- [8] P. Overholser, *Descendent Tropical Mirror Symmetry for \mathbb{P}^2* , preprint arxiv:math/1504.06138.
- [9] J. Rau, *Tropical intersection theory and gravitational descendants*, PhD thesis, TU Kaiserslautern, 2009.
- [10] J. Rau, *Intersections on tropical moduli spaces*, to appear in Rocky Mountain J. Math., 2015, preprint arxiv:math/0812.3678.

- [11] C. Torchiani, *Enumerative geometry of rational and elliptic tropical curves in \mathbb{R}^m* , PhD thesis, TU Kaiserslautern, 2014.

Falko Gauss, Lehrstuhl für Mathematik VI, Institut für Mathematik, Universität Mannheim, A5, 6, 68131, Mannheim, Germany
E-mail address: gauss@math.uni-mannheim.de