

Throughout this exercise sheet  $K = \mathbb{C}$  is the field of complex numbers,  $(G, \cdot)$  is a finite group, and  $V$  a finite-dimensional  $\mathbb{C}$ -vector space.

**EXERCISE 19**

Prove the following assertions:

(a) If  $G$  is a non-abelian simple group (or more generally if  $G$  is perfect, i.e.  $G = [G, G]$ ), then the image  $\rho(G)$  of any representation  $\rho : G \rightarrow GL(V)$  is a subgroup of  $SL(V)$ .

(b) No simple group  $G$  has an irreducible character of degree 2.

Assume that  $G$  is simple and  $\rho : G \rightarrow GL_2(\mathbb{C})$  is an irreducible matrix representation of  $G$  with character  $\chi$  and proceed as follows:

1. Prove that  $\rho$  is faithful and  $G$  is non-abelian.
2. Determine the determinant  $\det_\rho$  of  $\rho$ .
3. Prove that  $|G|$  is even and  $G$  admits an element  $x$  of order 2.
4. Prove that  $\langle x \rangle \triangleleft G$  and conclude that assertion (b) holds.  
(Use the diagonalisation theorem and steps 1., 2. and 3.)

**EXERCISE 20**

Let  $G$  be a finite group of odd order and, as usual, let  $r$  denote the number of conjugacy classes of  $G$ . Use character theory to prove that

$$r \equiv |G| \pmod{16}.$$

[Hint: Label the set  $\text{Irr}(G)$  of irreducible characters taking dual characters into account.]

**EXERCISE 21**

Prove that if  $\chi \in \text{Irr}(G)$ , then  $Z(G) \leq Z(\chi)$  and deduce that  $\bigcap_{\chi \in \text{Irr}(G)} Z(\chi) = Z(G)$ .

**EXERCISE 22 (Exercise to hand in / 8 points)**

(a) Let  $\rho : G \rightarrow GL(V)$  be an irreducible faithful  $\mathbb{C}$ -representation with character  $\chi$ . Let  $m \in \mathbb{Z}_{\geq 1}$  and let  $\rho^{\otimes m} := \rho \otimes \cdots \otimes \rho : G^m \rightarrow GL(V^{\otimes m})$  be the  $m$ -fold tensor product of  $\rho$  with itself.

- (i) Prove that  $Z(\chi) = Z(G)$ ;
- (ii) Prove that  $H := \{(z_1, \dots, z_m) \in Z(G^m) \mid z_1 \cdots z_m = 1\}$  is a normal subgroup of  $G^m$  such that  $|H| = |Z(G)|^{m-1}$  and  $H \leq \ker(\rho^{\otimes m})$ .
- (iii) Prove that  $\rho^{\otimes m}$  induces an irreducible  $\mathbb{C}$ -representation of  $G^m/H$  and deduce that  $\chi(1)^m \mid \frac{|G|^m}{|Z(G)|^{m-1}}$ .
- (iv) Set  $\alpha := \frac{\chi(1)}{\gcd(\chi(1), |G:Z(G)|)}$  and prove that  $\alpha^m \leq |Z(G)|$ .
- (v) Use (iv) to determine  $\alpha$  and deduce that  $\chi(1) \mid |G:Z(G)|$ .

(b) Deduce from (a) that  $\chi(1) \mid |G:Z(\chi)|$  for every irreducible character  $\chi \in \text{Irr}(G)$ .  
(Hint: mod out by the kernel of  $\chi$ .)