

Computer Algebra

Winter Semester 2013 - Problem Set 1

Due October 31, 2013, 2:00 pm

Problem 1. Let P be a totally ordered monoid and R be a P -filtered ring.

- (a) Let $0 \rightarrow M' \xrightarrow{\varphi} M \xrightarrow{\psi} M'' \rightarrow 0$ be a short exact sequence of R -modules and F_\bullet a P -filtration on M . Define by $F_p M' := \varphi^{-1} F_p M$ and $F_p M'' := \psi F_p M$ filtrations F_\bullet on M' and M'' . Show that the above sequence is strict with respect to F_\bullet .
- (b) Let $\varphi : M \rightarrow N$ be a P -filtered homomorphism, where M and N are P -filtered R -modules with filtrations F_\bullet and G_\bullet , respectively. Under a suitable hypothesis on F_\bullet and G_\bullet show that the following are equivalent:
- φ is strict with respect to F_\bullet and G_\bullet ,
 - $F_\bullet \varphi(M) = G_\bullet \varphi(M)$ (where the filtrations are defined as in part (a)),
 - $\text{gr}_\bullet^F \varphi(M) = \text{gr}_\bullet^G \varphi(M)$.

Note that there is a map $F_\bullet \varphi(M) \xrightarrow{\text{id}} G_\bullet \varphi(M)$.

Problem 2. The matrix $A \in \text{GL}(n, \mathbb{R})$ defines a monomial ordering $>_A$ on $\text{Mon}(x_1, \dots, x_n)$ by setting

$$x^\alpha >_A x^\beta :\Leftrightarrow A\alpha > A\beta,$$

where $>$ on the right-hand side is the lexicographical ordering on \mathbb{R}^n .

- Show that $>_A$ is indeed a monomial ordering on $\text{Mon}(x_1, \dots, x_n)$.
- Let $>$ be any monomial ordering on $\text{Mon}(x_1, \dots, x_n)$. Then there is a matrix $A \in \text{GL}(n, \mathbb{R})$ such that $>$ can be defined as $>_A$.

Problem 3. Consider a monomial ordering $>_1$ on $\text{Mon}(x_1, \dots, x_{n_1})$ and a monomial ordering $>_2$ on $\text{Mon}(y_1, \dots, y_{n_2})$. Then the product ordering or block ordering $>$, also denoted by $(>_1, >_2)$, on $\text{Mon}(x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2})$, is defined as

$$x^\alpha y^\beta > x^{\alpha'} y^{\beta'} :\Leftrightarrow x^\alpha >_1 x^{\alpha'} \text{ or } (x^\alpha = x^{\alpha'} \text{ and } y^\beta >_2 y^{\beta'}).$$

Given a vector $w = (w_1, \dots, w_n)$ of integers, we define the weighted degree of x^α by

$$w\text{-deg}(x^\alpha) := \langle w, \alpha \rangle := w_1 \alpha_1 + \dots + w_n \alpha_n,$$

that is, the variable x_i has degree w_i . For a polynomial $f = \sum_\alpha a_\alpha x^\alpha$, we define the weighted degree,

$$w\text{-deg}(f) := \max\{w\text{-deg}(x^\alpha) \mid a_\alpha \neq 0\}.$$

Using the weighted degree in the definition of $>_{dp}$, respectively $>_{ds}$ (cf. Example 1.2.8 in the SINGULAR book by Greuel, Pfister), with all $w_i > 0$, instead of the usual degree, we obtain the weighted reverse lexicographical ordering $>_{wp(w_1, \dots, w_n)}$, respectively the negative weighted reverse lexicographical ordering $>_{ws(w_1, \dots, w_n)}$. Now determine matrices $A \in \text{GL}(n, \mathbb{R})$ defining the orderings

- (a) $>_{ws(5,3,4)}$ on $\text{Mon}(x_1, x_2, x_3)$ with $n = 3$,
- (b) $(>_{dp}, >_{ls})$ on $\text{Mon}(x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2})$ with $n = n_1 + n_2$,
- (c) $(>_{ds}, >_{wp(7,1,9)})$ on $\text{Mon}(x_1, \dots, x_{n_1}, y_1, y_2, y_3)$ with $n = n_1 + 3$.

Problem 4. Write a SINGULAR procedure, having a list $P = ((g_1, h_1), \dots, (g_r, h_r))$ of pairs of polynomials, an ideal $I = \langle f_1, \dots, f_s \rangle$ and a polynomial f as input and returning the extended pair set $P = P \cup ((f, f_1), \dots, (f, f_s))$ as output.

Don't forget to add at least one example to your procedure.